arXiv:0709.4074v1 [hep-th]

Comment on “Hawking Radiation and Covariant Anomalies”

Shuang-Qing Wu* and Zhan-Yue Zhao

College of Physical Science and Technology, Central China Normal University, Wuhan, Hubei 430079, People’s Republic of China

We comment on the consistence of the epsilon anti-symmetric tensor adopted in [R. Banerjee and S. Kulkarni, arXiv:0707.2449] when it is generalized in the general case where $\sqrt{-g} \neq 1$. It is pointed out that the correct non-minimal consistent gauge and gravitational anomalies should by multiplied a factor $\sqrt{-g} \neq 1$. We also sketch the generalization of their work to the $\sqrt{-g} \neq 1$ case.

PACS numbers: 04.62.+v, 04.70.Dy, 11.30.-j

arXiv:0709.4074v1 [hep-th] 26 Sep 2007

Comment on “Hawking Radiation and Covariant Anomalies” In a recent paper [1], Banerjee and Kulkarni suggested that it is conceptually clean and economical to use only covariant gauge and gravitational anomalies to derive Hawking radiation from charged black holes [2]. Here we point out that while their work is self-consistent in the case of the metric determinant $\sqrt{-g} = \sqrt{-g_{tt}g_{rr}} = 1$, a potential ambiguity will arise in the definition of the epsilon anti-symmetric tensor when $\sqrt{-g} \neq 1$. We then remove this vagueness and present the correct expressions for the non-minimal consistent gauge and gravitational anomalies. Subsequently, we sketch the generalization of their work to derive the charge flux and energy-momentum flow in the most general diagonal metric case where $\sqrt{-g} \neq 1$.

Without loss of generality, let’s consider the most general two-dimensional non-extremal metric given by

$$ds^2 = f(r)dt^2 - h(r)^{-1}dr^2 \,. \qquad (1)$$

Here we assume $f(r_+) = h(r_+) = 0$ and $\sqrt{-g} = \sqrt{f/h}$ is regular at the horizon $r = r_+$. Then we define the two dimensional antisymmetric tensors, $\varepsilon^{\mu\nu} = \epsilon^{\mu\nu}/\sqrt{-g}$ and $\varepsilon_{\mu\nu} = \sqrt{-g}\epsilon_{\mu\nu}$ for the upper and the lower cases, respectively, together with $\epsilon^{01} = \epsilon_{10} = 1$. In terms of the usual Pauli matrices σ_i’s, the γ-matrices and γ_5-matrix are explicitly given by, $\gamma^0 = \sigma_2/\sqrt{f}$, $\gamma^1 = i\sqrt{h}\sigma_1$, $\gamma_0 = \sqrt{f}\sigma_2$, $\gamma_1 = -i\sigma_1/\sqrt{h}$, and $\gamma_5 \equiv \gamma_0\gamma_1/\sqrt{-g} = -\sigma_3$.

Using the property of γ-matrices, $\gamma_5\gamma^\mu = -\varepsilon^{\mu\nu}\gamma_\nu$, it is easy to check that the chiral current

$$\mathcal{J}^\mu = \bar{\psi}\Big(\frac{1 \pm \gamma_5}{2}\Big)\gamma^\mu\psi = \bar{\psi}\Big(\frac{g^{\mu\nu} \mp \varepsilon^{\mu\nu}}{2}\Big)\gamma_\nu\psi \,, \qquad (2)$$

satisfies a constraint condition, $\varepsilon_{\alpha\mu}\mathcal{J}^\mu = \mp\mathcal{J}_\alpha$.

The minimal form of the consistent gauge anomaly is

$$\nabla_\mu J^\mu = \pm\frac{e^2}{4\pi\sqrt{-g}}\epsilon^{\alpha\beta}\partial_\alpha A_\beta = \pm\frac{e^2}{4\pi}\varepsilon^{\alpha\beta}\partial_\alpha A_\beta \,, \qquad (3)$$

where $+(-)$ corresponds to left(right)-handed fields, respectively. Adding a local polynomial to the consistent current J^μ, then the current $\bar{J}^\mu = J^\mu + e^2A^\mu/(4\pi)$ is still consistent and should obey the anomalous equation

$$\nabla_\mu\bar{J}^\mu = \pm\frac{e^2}{4\pi\sqrt{-g}}\partial_\alpha\big[\sqrt{-g}\big(\varepsilon^{\alpha\beta} \pm g^{\alpha\beta}\big)A_\beta\big] \,. \qquad (4)$$

This is a non-minimal form for the consistent anomaly including the normal parity terms. Both consistent anomalies will obey the same equation once if the Lorentz gauge condition $\nabla_\mu A^\mu = 0$ is imposed. The new consistent current can be further modified, by adding a local polynomial, to define a covariant current

$$\tilde{J}^\mu = \bar{J}^\mu \mp \frac{e^2}{4\pi}A_\nu\big(\varepsilon^{\nu\mu} \pm g^{\nu\mu}\big) \,, \qquad (5)$$

which yields the gauge covariant anomaly

$$\nabla_\mu\tilde{J}^\mu = \pm\frac{e^2}{4\pi\sqrt{-g}}\epsilon^{\alpha\beta}F_{\alpha\beta} = \pm\frac{e^2}{4\pi}\varepsilon^{\alpha\beta}F_{\alpha\beta} \,. \qquad (6)$$

The minimal form of the consistent gravitational anomaly, for right handed fields, is

$$\nabla_\mu T^\mu_{\ \nu} = \frac{\epsilon^{\beta\delta}}{96\pi\sqrt{-g}}\partial_\delta\partial_\alpha\Gamma^\alpha_{\nu\beta} = \frac{\varepsilon^{\beta\delta}}{96\pi}\partial_\delta\partial_\alpha\Gamma^\alpha_{\nu\beta} \,. \qquad (7)$$

Including the normal parity terms, the non-minimal form for the consistent gravitational anomaly should read

$$\nabla_\mu\bar{T}^\mu_{\ \nu} = \frac{1}{96\pi\sqrt{-g}}\partial_\delta\partial_\alpha\Big[\sqrt{-g}\big(\varepsilon^{\beta\delta} + g^{\beta\delta}\big)\Gamma^\alpha_{\nu\beta}\Big] \,. \qquad (8)$$

Special care should be taken for the anti-symmetric tensor $\hat{\epsilon}_{\mu\nu}$ in the covariant gravitational anomaly

$$\nabla_\mu\tilde{T}^\mu_{\ \nu} = \frac{1}{96\pi\sqrt{-g}}\hat{\epsilon}_{\mu\nu}\partial^\mu R \,, \qquad (9)$$

where $\hat{\epsilon}_{\mu\nu} \equiv g_{\mu\alpha}g_{\nu\beta}\epsilon^{\alpha\beta} = \sqrt{-g}\varepsilon_{\mu\nu} = -g\epsilon_{\mu\nu}$ should be understood, in accordance with the conventions adopted in [2, 3].

Finally, the classical energy momentum tensor is traceless, due to an identity $T^\lambda_{\ \nu} = \varepsilon^{\lambda\mu}T_{\mu\nu}$, following from the chirality constraint.

The derivation of Hawking flux via covariant anomalies is completely parallel with the $\sqrt{-g} = 1$ case except some modifications as follows. All $\tilde{J}^r$ and $\tilde{T}^r{}_t$ appearing in [1] should be replaced by $\sqrt{-g}\tilde{J}^r$ and $\sqrt{-g}\tilde{T}^r{}_t$, respectively. Accordingly, the operator ∇_μ in Eqs. (22) and (31) should be multiplied by the factor $\sqrt{-g}$ also.

The surface gravity calculated by this method is

$$\kappa = \sqrt{-48\pi \tilde{N}^r{}_t(r_+)}\,, \tag{10}$$

$$\tilde{N}^r{}_t = \frac{1}{96\pi}\Big(hf'' + \frac{1}{2}h'f' - \frac{h}{f}f'^2\Big)\,. \tag{11}$$

Since $f(r_+) = h(r_+) = 0$, we find that $\tilde{N}^r{}_t(r_+) = \frac{-1}{192\pi}f'(r_+)h'(r_+)$, and so $\kappa = \frac{1}{2}\sqrt{f'(r_+)h'(r_+)}$, in agreement with that obtained in [3].

This work is partially supported by the NSFC under Grant No. 10675051.

Shuang-Qing Wu and Zhan-Yue Zhao

College of Physical Science and Technology, Central China Normal University, Wuhan, Hubei 430079, People's Republic of China

PACS numbers: 04.62.+v, 04.70.Dy, 11.30.-j

---

* Electronic address: sqwu@phy.ccnu.edu.cn

[1] R. Banerjee and S. Kulkarni, arXiv:0707.2449 [hep-th].

[2] S. Iso, H. Umetsu, and F. Wilczek, Phys. Rev. Lett. **96**, 151302 (2006).

[3] S.Q. Wu and J.J. Peng, Class. Quantum Grav. **24** (in press), arXiv:0706.0983 [hep-th].